

AD-A089 068

TEXAS UNIV AT AUSTIN DEPT OF CHEMISTRY
THE HETEROGENEOUS PHOTOCATALYTIC OXIDATION OF
AUG 80 I IZUMI, W W DUNN, K O WILBOURN
TR-10


F/G 7/3
HYDROCARBONS ON P-ETC(U)
N00014-78-C-0592
NL

UNCLASSIFIED

1 of 1
AUG 80



END
DATE
FILMED
10-80
DTIC



MICROCOPY RESOLUTION TEST CHART

NATIONAL BUREAU OF STANDARDS-1963-A

5L. **LEVEL** *IV*

(12)

OFFICE OF NAVAL RESEARCH

Contract N00014-78-C-0592

Task No. NR 051-693

TECHNICAL REPORT No. 10

AD A089068

The Heterogeneous Photocatalytic Oxidation
of Hydrocarbons on Platinized TiO_2 Powders

by

Ikuichiro Izumi, Wendell W. Dunn, Keith O. Wilbourn,
Fu-Ren F. Fan, and Allen J. Bard

Prepared for Publication

in the

Journal of ~~the American Chemical Society~~
Physical Chemistry

DTIC
ELECTE
SEP 11 1980
S D

The University of Texas at Austin
Department of Chemistry
Austin, Texas 78712

August 28, 1980

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited.

DDC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER (14) TR-10	2. GOVT ACCESSION NO. AD-A089 068	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) THE HETEROGENEOUS PHOTOCATALYTIC OXIDATION OF HYDROCARBONS ON PLATINIZED TiO_2 POWDERS.		5. TYPE OF REPORT & PERIOD COVERED 1 Sept. 1979 - 31 Aug. 1980
7. AUTHOR(s) (10) Ikuichiro Izumi, Wendell W. Dunn, Keith O. Wilbourn, Fu-Ren F. Fan, Allen J. Bard		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Texas at Austin Austin, TX 78712		8. CONTRACT OR GRANT NUMBER(s) (15) N00014-78-C-0592
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research 800 N. Quincy Arlington, VA 22217 (12) 222		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (11) 28 Aug 80
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (9) Technical Rept. 7 Sep 77-31 Aug 80		12. REPORT DATE August 28, 1980
		13. NUMBER OF PAGES 23
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication in the Journal of <i>Physical Chemistry</i>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) photocatalytic hydrocarbons		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The photodecomposition of hydrocarbons in oxygen-containing solutions at platinized TiO_2 yields predominantly CO_2 as the reaction product, with inter- mediate production of hydroxylated compounds. A mechanism for the reaction based on photogeneration of hydroxyl radicals at the TiO_2 surface is proposed.		

The Heterogeneous Photocatalytic Oxidation of Hydrocarbons on Platinized
TiO₂ Powders.

Ikuichiro Izumi, Wendell W. Dunn, Keith O. Wilbourn, Fu-Ren F. Fan, and
Allen J. Bard*

Department of Chemistry, The University of Texas at Austin, Austin, Texas 78712

(ABSTRACT)

The photodecomposition of hydrocarbons in oxygen-containing solutions at platinized TiO₂ yields predominantly CO₂ as the reaction product, with intermediate production of hydroxylated compounds. A mechanism for the reaction based on photogeneration of hydroxyl radicals at the TiO₂ surface is proposed.

(End of Abstract)

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or special
A	

Introduction

Recent investigations from this laboratory have described the application of platinized titanium dioxide powders (Pt/TiO_2) to heterogeneous photocatalytic and photosynthetic processes, such as the photo-Kolbe reaction, in which aliphatic mono-¹ and di-² carboxylic acids are decomposed to give the corresponding alkane as the main product, as well as other reactions³. These investigations have clearly demonstrated the usefulness of Pt/TiO_2 in photoelectrochemical experiments and the potential for the accomplishing reactions requiring highly oxidizing conditions. In previous works we found that the decarboxylation of benzoic acid at Pt/TiO_2 involved the preceding hydroxylation of the benzene ring to form hydroxylated benzoates as reaction intermediates.² However, the main product of this reaction was CO_2 , suggesting that photodecomposition of hydrocarbon intermediates was possible. Thus in this paper we extend these studies to the photocatalyzed decomposition of hydrocarbons at Pt/TiO_2 . Phenol is shown to be a reaction intermediate for the photodecomposition of benzene, with the final product, CO_2 . This unusual and efficient photocatalyzed breakdown of benzene at room temperature to yield CO_2 thus provides a possible pathway in the decarboxylation mechanism proposed for the photooxidation of benzoic acid.² We also describe the photodecomposition of several aliphatic hydrocarbons, in which alcohols have been detected as intermediates.

Experimental Section

Materials. Benzene [reagent grade, Matheson, Coleman and Bell (MCB)], hexane (MCB, 99⁺ mol %), cyclohexane (spectrophotometric grade, MCB), heptane (reagent grade, Eastman), nonane and decane (Fisher Certified), barium hydroxide (reagent grade, MCB) and phenol (reagent grade, Fisher) were used without further purification. Decanol, nonanol, heptanol, hexanol, and cyclohexanol

(reagent grade, Eastman) were used as received for standards in GC-Mass spectroscopic analysis. The kerosene was extracted repeatedly with distilled water prior to illumination to simplify subsequent analysis of the aqueous phase. All solvents and other chemicals were reagent or spectrophotometric grade and were used without further purification. The platinized TiO_2 powders were prepared by photodecomposition of hexachloroplatinic acid onto TiO_2 powders (reagent grade, MCB, 125-250 μm) and contained 10 % platinum by weight.⁴

Apparatus. The irradiation source for the photodecomposition of aliphatic hydrocarbons was an Atlas Weatherometer, Model 6000, equipped with a 6000 W Xe lamp. The described experiments were performed at an unfocused power output of 5000 Watts, which corresponds to a total radiant intensity of $\sim 35 \text{ mW/cm}^2$ at the location of the reaction cell. The Xe lamp was jacketed with a water cooling cell to remove infrared irradiation. A constant reaction temperature was maintained with a large water bath fitted with a coil for water cooling. A 2500 W Xe lamp (Model UF 30 KK, Christie Electric Corp., Los Angeles, Calif.) operated at 1600 W, was used as the light source for all preparative runs in the photodecomposition of benzene. The reaction cell with a flat window for irradiation and the water bath were both of Pyrex. A 450 W Xe lamp with Model 6242 power supply (Oriel Corp., Stamford, Conn.) served in the photoelectrochemical measurements. Mass or GC-Mass spectra of reaction gases were obtained with an automated Gas Chromatograph/EI-CI Mass Spectrometer System (Finnigan, Model 2000). UV-visible spectra were recorded with a Cary Model 14 spectrophotometer. Electrochemical experiments were performed using a PAR Model 173 Potentiostat (Princeton Applied Research, Princeton, NJ) and a PAR Model 173 Universal Programmer.

Product Analysis. (a) Aliphatic Hydrocarbons: In a typical experiment, 200 mg of the photocatalyst powder was suspended by magnetic stirring in a two-phase system composed of 5 mL water and 5 mL of the hydrocarbon. The reaction cell was maintained at 36 ± 1 °C. The sweep gas, N_2 or O_2 , was first passed through $Ba(OH)_2$ to remove any traces of CO_2 in the cylinder gas and then into the reaction cell. The gas was then passed through a large volume cell cooled at -56 °C to condense volatile organic chemicals, and then into a $Ba(OH)_2$ solution for determination of CO_2 as precipitated $BaCO_3$. Solar experiments were conducted in a Pyrex flask with a liquid surface area of approximately 700 cm^2 using 1.0 g of the photocatalyst. The condensed volatile organic chemicals were analyzed by a GC-Mass spectrometer. For hydrocarbon analysis, the columns used were a 6 m capillary of OVI and a 36 cm OVI column. Relative concentrations of trace impurities were compared by integration of ion current peaks and calibrated with standard samples. Investigation of the more polar constituents was performed with a Carbowax column. Identification of intermediate products was based on comparison of retention times and mass spectra of sample and standards. A detection limit of 2 ppm (equivalent to $\sim 12\text{ }\mu\text{M}$) was established for heptanol under these conditions.

The gas products not condensed at -56 °C were collected in a gas buret and were analyzed by a mass or GC-Mass spectrometer.

(b) Fenton's Reagent Reaction of Heptane: 5 mL heptane suspended in 100 mL 0.64 M $FeSO_4$, was purged with N_2 to remove CO_2 . 5 mL H_2O_2 was added while the system was swept with N_2 . After 12 hours, an additional 3 mL of H_2O_2 was added to ensure complete reaction. The evolved CO_2 was precipitated as $BaCO_3$ from a saturated $Ba(OH)_2$ solution.

(c) Benzene: In a Pyrex cell, a stirred suspension of 100 mg of Pt/TiO₂ in an emulsion of 5 mL benzene and 20 mL distilled water held at 27 ± 1 °C was illuminated under oxygen with white light from a 2500 W Xe lamp (operated at 1600 W). The gases produced during the irradiation were collected in a mercury-containing gas volumetric apparatus, as reported previously.¹ After termination of photolysis, the mixture in the gas buret was transferred to an evacuated gas sample cell which attached to the mass spectrometer for gas analysis. The average rate of CO₂ evolution was determined gravimetrically in another preparative run. The reaction gases were swept out of the reaction vessel with a stream of oxygen and were bubbled through a saturated solution of Ba(OH)₂ in 0.2 M sodium hydroxide. The precipitated BaCO₃ was filtered off, washed with distilled water, dried at 120 °C, and weighed. The colorless reaction mixture after filtering off the photocatalyst powders was analyzed by UV-visible absorption.

Results

Photocatalytic Decomposition of Benzene on Platinized TiO₂ Powders.

Illumination of suspensions of platinized TiO₂ powders in aqueous benzene mixtures in the presence of oxygen produced observable evolution of gaseous CO₂. After illumination, the gaseous products were analyzed by mass spectrometry. The mass spectra [m/e (relative intensity)] consisted of signals of CO₂, 44 (44.9 %), 16 (11.5 %) with background signals due to O₂, 32 (100 %); N₂, 28 (28 %); benzene, 78 (12 %); and water, 18 (5 %). Thus the only significant gaseous product of photooxidation of benzene in water was CO₂. The yield of CO₂ determined gravimetrically in another series of experiments is shown in Table I. The UV spectrum of the colorless reaction mixture showed absorption maxima at 2390, 2720, and 2780 Å with a shoulder at 2670 Å. The product was identified as phenol by comparing its UV spectrum with that of a standard

phenol solution. The concentration of phenol given in Table I was determined by UV spectroscopy. The results of some control experiments are also summarized in Table I. As shown in experiment 4, the photodecomposition of benzene requires the presence of catalyst. Platinized anatase has a higher efficiency than plain anatase powder (see experiments 2 and 7). The results shown in experiments 3 and 6 show that the presence of oxygen promotes the photodecomposition of benzene. Particularly interesting, the addition of water, compared to a system containing only benzene, promotes the photodecomposition of benzene (see experiments 2 and 5). As shown in experiments 1 and 2, substantial amounts of phenol were isolated as an intermediate product. These results suggest that the photooxidation of benzene on platinized TiO_2 powders suspended in H_2O probably proceeds via the hydroxylation of benzene to form phenols as the intermediate.^{5,6} Moreover, the results are different from those for the electrolysis of aqueous benzene solutions in which p-benzoquinone is the predominant product.⁷⁻⁹ No p-benzoquinone or p-hydroquinone were obtained in the present experiments.

Photocatalytic Decomposition of Aliphatic Hydrocarbons on Pt/ TiO_2 Powders.

Illumination of a suspension of Pt/ TiO_2 in aqueous aliphatic hydrocarbon mixtures in the presence of oxygen also leads to the observable evolution of CO_2 . The yields of CO_2 , which was the only significant gaseous product (not condensed at -56°C), are shown in Table II.

Extensive analysis for intermediates revealed a detectable level of only the alcohols (see Table II), except in the case of decane, where trace amounts of 2-, 4-, and 5-decanone were identified. The concentrations of lower molecular weight hydrocarbon impurities remained essentially unchanged before and after reaction, demonstrating that the cleavage of the alkanes does not result in the formation of shorter-chain hydrocarbons.

The reduction in the rate of photocatalytic oxidation with increasing molecular weight is coincident with an increase in the viscosity of the hydrocarbons. An exception to this trend is cyclohexane; however, in this case an emulsion of the three phases (photocatalyst powder, H_2O , and hydrocarbon) occurs. Addition of an emulsifying agent, IGEPAL-560 (nonionic nonyl phenoxy polyoxyethylene ethanol, GAF Corp.) to a suspension of decane was found to increase the rate of CO_2 production from 34.0 $\mu\text{mole/hr}$ to 75.0 $\mu\text{mole/hr}$, confirming that the enhanced rate for cyclohexane oxidation is probably due to improved phase mixing.


The reaction between heptane and Fenton's reagent, which produces hydroxyl radicals, also produced a considerable amount of CO_2 (Table II). This demonstrates the possibility that at least in the initial stages of the photodecomposition process, the attack of $\cdot OH$ on hydrocarbons and intermediates occurs.

Electrochemical and Photoelectrochemical Measurements. The photocatalytic activities of suspended platinized anatase powders were correlated with the behavior of TiO_2 single crystal electrodes in photoelectrochemical (PEC) measurements as shown in Figure 1. The current-potential behavior of a rutile single crystal electrode was examined in an aqueous solution containing 1 M $NaClO_4$ as the supporting electrolyte. In the dark, only a very small anodic current was observed on a TiO_2 electrode both in deoxygenated and in oxygen-saturated solutions, with or without benzene. In oxygen-saturated solution, however, the TiO_2 electrodes showed a reduction peak at -1.16 V vs. SCE (curve 2); this peak can be attributed to the reduction of oxygen, since it disappeared when the solution was thoroughly deoxygenated (curve 1). These voltammetric curves were essentially unaffected by saturating the solution with benzene (curves 5 and 6). Under illumination, an appreciable anodic

current was observed. The on-set photopotential, V_{on} , was essentially unaffected by the presence of oxygen or benzene, and was about -0.35 V vs. SCE for all cases. The presence of oxygen did not change the magnitude of the photocurrent, however, the presence of benzene caused a slight reduction in the photocurrent (curves 7 and 8), probably due to the absorption of some of the incident light by benzene and phenols generated. On the other hand, proton reduction on a platinum electrode in the presence of benzene occurred at potentials more negative than -0.4 V vs SCE (curve 9). However, oxygen reduction on platinum occurred at potentials much more positive than either the on-set photopotential or the reduction potential of oxygen on TiO_2 (curve 10). Thus, the presence of platinum on the TiO_2 powder could promote the photocatalytic activity of TiO_2 compared to TiO_2 alone. Also, the photocatalytic activity of platinized TiO_2 powder would be substantially enhanced by the presence of oxygen. These results at a rutile electrode are thus qualitatively consistent with the powder results. The role of oxygen in PEC and the accompanying photocatalytic reactions on TiO_2 have also been demonstrated by Korsunovskii.¹⁰

Discussion

In the presence of oxygen, several n-type semiconductor powders have been shown to behave as photocatalysts and promote the oxidation of substrates. Irradiation of a semiconductor with light of energy higher than the band gap results in the creation of holes in the valence band and electrons in the conduction band of the semiconductor. These charge carriers can recombine or the holes can be scavenged by oxidizable species (for example, H_2O , H_2O_2 , or hydrocarbons (RH)), and electrons by reducible species (for example, O_2 or H^+) in the solution.



The electrochemical and photoelectrochemical measurements suggest that the reduction of H^+ on Pt and the reduction of O_2 on TiO_2 do not occur at an appreciable rate at the potentials where significant anodic photocurrent on TiO_2 is observed. However, the reduction of O_2 on Pt occurs at these potentials and this represents a viable half-reaction at the photocatalyst particle. The role of the Pt for this reaction is thus to provide a site for the more efficient utilization of the photogenerated electrons in the reduction of O_2 (e.g., eq. 4), in agreement with the finding that the photooxidation of hydrocarbons in deoxygenated solutions does not occur to an appreciable extent (Tables I and II). It is also possible that intermediates formed during the reduction of O_2 play an important role in the oxidation process, as suggested by reactions (4) - (6). Recent experiments involving spin trapping and electron spin resonance spectroscopic detection of intermediates formed during irradiation of Pt/TiO_2 in aqueous solutions have demonstrated the intermediacy of $\cdot\text{OH}$ and $\text{HO}_2\cdot$.¹⁷ The very slow rate of CO_2 production in pure benzene (see Exp. 5 in Table I), the lack of the intermediates other than hydroxylated compounds, and the dependence of the CO_2

evolution rate on the degree of phase mixing (Table II) suggest that the photogenerated holes are scavenged by water (eq. (2)) rather than by hydrocarbons (eq. (8)). These would form $\cdot\text{OH}$ or $\cdot\text{HO}_2$, and lead to the hydroxylation of hydrocarbons and eventually the complete oxidation. The mode of the complete decomposition of the hydrocarbons has not been established. However, the detection of phenol for the case of benzene and alcohols for the cases of aliphatic hydrocarbons suggests that attack by $\cdot\text{OH}$ or HO_2 radicals is a likely first step. The probable role of $\cdot\text{OH}$ in these reactions is also supported by the reaction of benzoic acid² and hydrocarbons with Fenton's reagent.

Thus the heterogeneous photooxidation of hydrocarbons in suspensions with Pt/TiO_2 can be explained by a mechanism similar to that invoked for benzoic acid, that is, the photooxidation proceeds via hydroxylated intermediates. In both experiments it is of importance that hydrocarbons, which are unattacked by most oxidizing agents, readily react with $\cdot\text{OH}$ to form the final product CO_2 . Since aliphatic hydrocarbons are themselves the products of the photo-Kolbe reaction,^{1,2} their isolation in experiments involving the photodecomposition of organic acids suggest that they are capable of escaping from the powder surface. They then are swept from the reaction medium or react with $\cdot\text{OH}$ less readily than the organic acid precursor. The destruction of hydrocarbons by the photocatalytic reaction described here might be useful in the treatment of waste streams or spills involving these materials.

Acknowledgment

The support of this research by the Educational Ministry of Japan (to I.I.), by the Office of Naval Research (to F.F.F.), and the National Science Foundation is gratefully acknowledged.

References

- (1) (a) B. Kraeutler and A. J. Bard, J. Am. Chem. Soc., **100**, 2239 (1978);
(b) B. Kraeutler and A. J. Bard, J. Am. Chem. Soc., **100**, 5985 (1978).
- (2) I. Izumi, Fu-Ren F. Fan and A. J. Bard, J. Phys. Chem. (submitted).
- (3) (a) H. Reiche and A. J. Bard, J. Am. Chem. Soc., **101**, 3127 (1979);
(b) S. F. Frank and A. J. Bard, J. Phys. Chem., **81**, 1484 (1977).
- (4) B. Kraeutler and A. J. Bard, J. Am. Chem. Soc., **100**, 4317 (1978).
- (5) (a) G. A. Korsunovskii, Zhur. Fiz. Khim., **31**, 2351 (1957); (b) N. A. Vysotskaya, Uspekhi Khim., **42**, 1848 (1973) (Russ. Chem. Rev., No. **10**, 1973).
- (6) N. L. Weinberg, Ed., "Technique of Electroorganic Synthesis", Wiley-Interscience, New York, N.Y., 1974, p. 251.
- (7) J. S. Clarke, R. E. Ehigamusue and A. T. Kuhn, J. Electroanal. Chem., **70**, 333 (1976).
- (8) S. Ito, K. Sasaki, Y. Murakami and H. Shiba, Denki Kagaku, **40**, 733 (1972).
- (9) (a) K. S. Udupa, G. S. Subramanian and H.V. K. Udupa, Pol. De. Sci., **9** (1961); (b) Y. Isomura, J. Chem. Soc. Japan, **62**, 1167 (1947).
- (10) G. A. Korsunouskii, Russ. J. Phys. Chem., **52**, 1314 (1978).
- (11) T. Freund, and W. P. Gomes, Catal. Rev., **3**, 1 (1969) and references therein.
- (12) A. J. Bard, J. Photochem., **10**, 59 (1979).
- (13) A. J. Bard, Science, **207**, 139 (1980), and references therein.
- (14) M. Formenti, and S. J. Teichner, Specialists Periodical Reports (The Chemical Society, London), **2**, 1979, Chapter 4.
- (15) S. N. Frank and A. J. Bard, J. Am. Chem. Soc., **99**, 4667 (1977).
- (16) S. N. Frank and A. J. Bard, J. Phys. Chem., **81**, 1484 (1977).
- (17) C. D. Jaeger and A. J. Bard, J. Phys. Chem., **83**, 3746 (1979).

TABLE I. Photocatalytic Oxidation of Benzene on Platinized Anatase or Anatase Powders.^a

<u>Exp. No.</u>	<u>Catalyst</u>	<u>Time of illumination hrs</u>	<u>Conditions</u>	<u>CO₂ generated mmol</u>	<u>Phenol generated mmol</u>
1	10% platinized anatase	4	illuminated, O ₂ saturated	0.13	0.10
2	10% platinized anatase	18.5	illuminated, O ₂ saturated	0.72	0.41
3	10% platinized anatase	40	illuminated, O ₂ saturated	1.85	---
4	none	72	illuminated, O ₂ saturated	0.03	---
5 ^b	10% platinized anatase	19	illuminated, O ₂	0.10	---
6	10% platinized anatase	43	illuminated, deoxygenated	0.15	---
7	anatase	14	illuminated, O ₂ saturated	0.38	---

^aIf not otherwise mentioned, the solution was composed of 5 mL of benzene suspended in 20 mL of H₂O. 100 mg of catalyst was employed.

^b25 mL of benzene was used as the reacting solution.

TABLE II. Photocatalytic Oxidation of Aliphatic Hydrocarbons on Platinized Anatase
Powders.

Alkane	Weight Powder (g)	Time of illumination (hr)	Rate CO ₂ production (μ mol/hr)	Rate of oxidation (μ mol/hr)	Alcohol generated (μ mol)
Hexane	0.199	24	53.7	8.95	
	0.200	12	30.9	5.2	0.55
	0.200	24	65.2	10.9	
Cyclohexane	0.200	24	86.4	14.4	6×10^{-5}
Heptane	0.200	48	60.4	8.6	0.650
	0.201	24	61.1	8.7	
	0.201	24	45.3	6.5	
	^a 0.0	18	14.4	2.1	
Nonane	0.200	24	41.4	4.6	0.220
	0.200	46	65.6	7.3	
	^b 1.430	24	54.6	6.1	
Decane	0.200	67	14.8	1.5	
	0.201	291	34.1	3.4	0.105
	^c 0.0	30	0.0	0.0	
	^d 0.200	24	0.0	0.0	
	^e 0.200	6	75.0	7.5	
Kerosene	0.200	72	17.2	---	---

^aPerformed with Fenton's Reagent, see exp. section.

^bSolar experiment.


^cControl experiment, no catalyst added.

^dN₂ substituted for O₂.

^eEmulsifying agent added (0.25 mL IGOPAL - 560)

FIGURE CAPTION

- Fig. 1** Voltammetric curves at platinum and single crystal rutile electrodes in solution containing 1 M NaClO_4 as the supporting electrolyte. Scan rate 100 mV/sec. Initial potential at the positive extremes.
- Curve 1: In the dark; on TiO_2 ; under N_2 ; solution without containing benzene.
- Curve 2: In the dark; on TiO_2 ; oxygen saturated; without benzene.
- Curve 3: Under illumination; on TiO_2 ; under N_2 ; without benzene.
- Curve 4: Under illumination; on TiO_2 ; under O_2 ; without benzene.
- Curve 5: In the dark; on TiO_2 ; under N_2 ; benzene saturated solution.
- Curve 6: In the dark; on TiO_2 ; oxygen and benzene saturated solution.
- Curve 7: Under irradiation; on TiO_2 ; under N_2 ; benzene saturated solution.
- Curve 8: Under irradiation; on TiO_2 ; oxygen and benzene saturated solution.
- Curve 9: On Pt; under N_2 ; benzene saturated.
- Curve 10: On Pt; oxygen and benzene saturated.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CND-AA-1P P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO9 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auborn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co, Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. H. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Larry F. Plew Naval Weapons Support Center Code 30736, Building 2905 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
S. Ruby DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton SO9 5NH England	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
1	Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1 Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, DC 20376
1	Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1 M. L. Robertson Manager, Electrochemical Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522
1	Dr. M. G. Sceats Department of Chemistry University of Rochester Rochester, New York 14627	1 Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720
1	Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1 Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709
1	Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1 Dr. Denton Elliott Air Force Office of Scientific Research Bldg. 104 Bolling AFB Washington, DC 20332
1	Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1
1	Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	